

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
19 February 2004 (19.02.2004)

PCT

(10) International Publication Number
WO 2004/014929 A1

(51) International Patent Classification⁷: C07H 15/18,
5/06, 23/00

ABBENANTE, Giovanni [AU/AU]; 53 Pringles Road,
SAMPSONVALE, Queensland 4520 (AU). LIU, Ligong
[AU/AU]; 28 Samara Street, SUNNYBANK, Queensland
4109 (AU).

(21) International Application Number:
PCT/AU2003/001008

(74) Agent: CULLEN & CO.; Level 26, 239 George Street,
BRISBANE, Queensland 4000 (AU).

(22) International Filing Date: 8 August 2003 (08.08.2003)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:
2002950657 8 August 2002 (08.08.2002) AU

(71) Applicant (*for all designated States except US*): AL-
CHEMIA PTY LTD [AU/AU]; 3 Hi-Tech Court, Brisbane
Technology Park, EIGHT MILE PLAINS, Queensland
4113 (AU).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): WEST, Michael,
Leo [AU/AU]; 364 Hemmant and Tingalpa Rd, HEM-
MANT, Queensland 4174 (AU). ANDREWS, Peter
[AU/AU]; 311 Swann Road, ST LUCIA, Queensland 4067
(AU). RAMSDALE, Tracie, Elizabeth [AU/AU]; 160
Hellawell Road, SUNNYBANK HILLS, Queensland 4109
(AU). MEUTERMANS, Wim [BE/AU]; 293 Birdwood
Terrace, TOOWONG, Queensland 4066 (AU). THANH
LE, Giang [AU/AU]; 38 Tarrant Street, MT GRAVATT,
Queensland 4122 (AU). CLARK, Chris [AU/AU]; 4/21
Waverly Road, TARINGA, Queensland 4068 (AU).


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- with amended claims

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DERIVATIVES OF MONOSACCHARIDES FOR DRUG DISCOVERY



(I)

(57) Abstract: New compounds and methods for the preparation of combinatorial libraries of potentially biologically active compounds are based on monosaccharides of formula (I) being a derivative of a furanose or pyranose form of a monosaccharide.

DERIVATIVES OF MONOSACCHARIDES FOR DRUG DISCOVERY

FIELD OF THE INVENTION

This invention relates to new compounds and methods for the preparation of combinatorial libraries of potentially biologically active compounds based on natural and unnatural monosaccharides.

These compounds are functionalized, with a view to varying lipid solubility, size, function and other properties, with the particular aim of discovering novel drug or drug-like compounds, or compounds with useful properties. The invention provides intermediates, processes and synthetic strategies for the solution or solid phase synthesis of monosaccharides, variously functionalised about the sugar ring, including the addition of aromaticity and charge, the addition of pharmacophoric groups and the placement of amino acid and peptide side chain units or isosteres thereof.

BACKGROUND OF THE INVENTION

In the field of drug discovery there is a constant need for novel scaffolds that enable the rational design of potentially bioactive molecules. Carbohydrates have recently come under scrutiny as offering a source of scaffolds that allow for a high degree of substitution, and offer access to both functional and structural diversity. The nature of monosaccharide molecules is such that there are numerous different stereoisomers available that can provide access to a greater degree of molecular space than do the scaffolds presently employed in drug discovery.

Carbohydrate monomers predominantly contain hydroxyl groups but also may contain other functionalities such as an amino and/or carboxylate function. In essence, the concepts involved in drug discovery through carbohydrate based molecular and structural diversity, are twofold: (1) The primary concept involves the exploitation of the high functional density found around the carbohydrate ring to display several different moieties of biological relevance. There is a dual significance to this substitution in that (i) the substituents relative position around the ring may be varied in relation to each other and, (ii) each individual moiety may be substituted for a class of such moieties and therefore themselves may be varied (by example: an arginine

mimetic may be substituted at position 1, 2, 3, 4 or 5 around a ring in relation to other peptidomimetics, by the same token the arginine mimetic may represent a class of different arginine bioisosteres which may all be similarly substituted).

(2) The second concept involves exploiting the structural diversity inherent in carbohydrate isomers. Each of the substituents around a carbohydrate ring may theoretically be presented in either an axial or equatorial configuration allowing access to hugely diverse molecular space. Many monosaccharides are naturally occurring, which aside from being useful in their own right, present themselves as cheap starting materials to access more exotic configurations.

There are other factors that promote carbohydrates as useful building blocks for drug discovery, for example the relative positions of the functional groups on the sugar rings are conveniently spaced such that they can effectively enable mimicry of (for example), peptide motifs such as peptidic turns and loops, as well as cyclic peptides.

The major difficulty encountered in attempts to employ monosaccharides as scaffolds, is associated with monosaccharide chemistry. In the past carbohydrate chemistry was considered arduous, protracted and not cost effective. Particularly, the degree of orthogonal protection group chemistry required to allow free access to any one of a monosaccharide's functional groups (usually five) was deemed too high to ever be effected in a commercially viable manner. As a corollary, the more easily effected peptide synthesis only requires a maximum three orthogonal protecting groups, additionally the conditions required for peptide synthesis are often milder, thus peptide synthesis has so far been able to be effected more easily than carbohydrate synthesis. Fortunately, recent developments in synthetic carbohydrate chemistry have begun to allow regular access to carbohydrates as molecular scaffolds. In a recent patent application (PCT AU00/00025) we disclosed a range of orthogonally protected building blocks suitable for oligosaccharide synthesis. The building blocks presented in this application are also suitable for use as intermediates in the synthesis of compounds of the present invention, and represent compounds and methods which define the state of the art.

A large number of Carbohydrate based templates and scaffolds has now been published in the scientific literature. A review of the major contributions by Gruner et. al., (Chem. Rev., 2002, 102, p491-514) highlights this activity. Within the general literature, there are two distinct types of carbohydrate templates (i) sugar amino acids and (ii) carbohydrate scaffolds.

Sugar amino acids are carbohydrates which contain both an amine function and a carboxylic acid function, and are used in place of amino acids in peptide type syntheses. The synthesis of monosaccharides for this purpose is exemplified by the work of Fleet (Tetrahedron, 1996, 52, p10711; Tetrahedron Assym., 1996, 7, p387; Tetrahedron Assym., 1996, 7, p157) and Le Merrer (Tet. Lett., 1995, 36, p6887) for furanoid sugars, and by Dondoni (J.Org.Chem., 1994, 59, p6404), Vogel (J. Carbohyd. Chem., 1994, 13, p37) and Kessler (see chem rev. above) for pyranoid sugars.

Sugar amino acids have been used in peptide synthesis, and in the formation of linear oligomers for various biological purposes (see chem reviews above). Importantly, all of these compounds contain an amino function and a carboxylate function directly attached to the carbohydrate ring, and these functional groups are involved in amide bond forming processes which is the central concept in their use. The compounds of this type are distinctly different from the compounds of the present invention.

Carbohydrate scaffolds have also received considerable attention in the scientific literature, at least by way of desideratum. In concept, these compounds provide a chiral scaffold on which pharmaceutically active moieties are presented. This is the field of the present invention which adds to and is distinct from the state of the art.

The use of carbohydrates as scaffolds was promulgated by Hirschmann and co workers (Hirschmann et. al., J. Am. Chem. Soc., 114, 9217-9218, 1992) who employed this concept to develop a potent NK-1 receptor antagonist (Hirschmann et. al., J. Am. Chem. Soc., 115, 12550-12568, 1993), (Hirschmann et. al., J. Med. Chem., 39, 2441-2448, 1996). The fundamentals of this work have also been patented by Hirschmann et. al. (PCT/US1994/012233).

In a similar manner, Papageorgiou et al, have applied the concept to furanoid structures, developing weak somatostatin inhibitors in the process (Papageorgiou et. al., Bioorg. Med. Chem. Lett., 2, 135-140, 1992).

Weak inhibitors of integrin receptors and endothelin receptors have also been developed by applying this concept (Nicolaou, K.C., et. al, Tetrahedron, 1997, 53, p8751; Moitessier, N., et. al., Lett. Pep. Sci., 1998, 5, p75; Moitessier, N., et. al., Bioorg. Med. Chem., 2001, 9, p511.).

A number of other research groups have developed libraries of compounds based on this scaffold principle, and these groups are referred to in Gruner's review (vide supra). Despite the plethora of work to date, the compounds disclosed above have three common features which distinguish them from the current work: (i) all of the substituents are attached to the scaffold through an oxygen linkage, (ii) the anomeric position is always an O glycoside, and (iii) all of the available hydroxyl positions are substituted.

These features, when taken together, place significant limitations on the utility of the compounds. For example, ether linkages provide considerable rotational freedom and it is generally accepted that rotational freedom often results in diminished biological activity (Murphy et. al., J. Org. Chem., 68, 5692-5704, 2003). To this end, the present invention is directed to carbohydrate templates which have one or two amines directly attached to the carbohydrate ring, allowing the introduction of, for example, amide linked, sulfonamide linked, urea linked and carbamoyl linked moieties with significantly reduced rotational freedom and often better physical properties.

In a similar manner, the requisite for all of the positions to be substituted can lead to compounds of higher lipophilicity, higher molecular weight and lower solubility without imparting greater biological activity. In the present invention we disclose compounds with one or two hydroxyl positions unsubstituted, allowing generally improved solubility characteristics and lower molecular weights that would be expected for the corresponding fully substituted molecules.

These two features represent significant improvements over compounds described in the literature and are the result of considerable new method developments by the inventors.

Of all the carbohydrate scaffold work reported in the scientific and patent literature to date, we have found few examples of amine containing scaffolds outside the sugar amino acid class. Kunz et. al. (WO 99/07718) have claimed 2-deoxy 2-amino sugars as scaffolds for drug discovery. This citation does not teach or exemplify a compound with an amine group directly attached to the ring in the two position or any other position.

The disclosures in Kunz's relate specifically to the use of glucose, galactose and mannose as scaffolds and the methods described are not generally applicable to other monosaccharide scaffolds. In contrast, the compounds of the present invention are all O glycosides which are further limited by a narrow range of unsubstituted substituents dictated by the low reactivity of the sugar hydroxyls under the synthetic conditions disclosed. It is apparent that this technology displays significant disadvantages to the present invention; the efficiencies of conversion, the range of potential substituents, the various inversion chemistries that introduce both alternate oxy and amino stereochemical orientations, and the versatile alkylative chemistries of the present invention represent significant improvements over the methods of Kunz's application. Particularly, the present invention provides stereoisomers of monosaccharides that have a nitrogen or a carbon atom attached to the ring in positions 3,4,5 and 6 of a monosaccharide or tetrahydrofurano/pyrano ring system. Of particular interest to the medicinal chemist is the inclusion of linking functionalities that are likely to be stable to physiological conditions thus allowing the drug to reach the desired target intact, or in an active form.

Despite the general paucity of amine containing carbohydrate scaffolds in the literature, there are many examples of monosaccharide building blocks and protected aminosugars employed for oligosaccharide synthesis. By way of example, US 4818816 discloses a compound 1-methyl-2-carbobenzyloxy,3-benzyl glucosamine, a monosaccharide building block used in the synthesis of synthetic heparinoid oligomers. The compounds of the

present invention represent a significant departure from the simple building block type aminosugars, both in the diversity and complexity which is achievable. In order, to further distinguish the compounds of the present invention from the prior art, the use of standard amine protecting groups in carbohydrate synthesis is specifically excluded.

Sabesan (US patent 5,220,008) discloses a series of higher oligosaccharides as inhibitors on influenza. Within the claims of this patent, a partially protected monosaccharide (structure IV) is also disclosed. The compounds of this structure are protected monosaccharides for oligosaccharide synthesis which are known in the art and do not represent compounds for drug discovery.


Similarly, Alchemia Pty Ltd has disclosed in PCT/AU01/01307 building blocks, methods of syntheses, and final products relating to the employment of monosaccharide compounds as drug like molecules. The compounds of PCT/AU01/01307 are specifically directed at inhibitors of the muramyl cascade of enzymes and are hereby excluded from specification by the incorporation of this reference. A number of other publications relating to muramyl type compounds have appeared in the literature. Liu et. al. (Biorg. Med Chem Lett., 10, 2000, 1361-1363) present a series of compounds containing a benzyl glycoside at the anomeric position, an acetate at C-2 and a peptide homologated lactate at C-3 of a glucosamine scaffold. These compounds and those disclosed by Xiao (Peptides: Biol and Chem., Proc. 5th Int. Chinese Peptide Symp., 1998 CA: 134:178795) represent compounds and methods which help define the art of carbohydrate chemistry but are not directly relevant to the present invention.

It will be clearly understood that, if a prior art publication is referred to herein, this reference does not constitute an admission that the publication forms part of the common general knowledge in the art in Australia or in any other country.

30

OBJECT OF THE INVENTION

In a first aspect, the invention comprises a compound of formula I being a derivative of a furanose or pyranose form of a monosaccharide,



formula I

Wherein, n is 0 or 1;

5

R1 is XR wherein,

X is selected from O; S; S=O and SO₂,

R is selected from the group consisting of C1 to C9 alkyl, C1 to C15 alkenyl, C1 to C15 alkynyl, C1 to C15 heteroalkyl, C6 to C15 aryl, C6 to C15 heteroaryl, C6 to C15 arylalkyl or C6 to C15 heteroarylalkyl which is optionally substituted,
10 cyclic or acyclic, branched and/or linear,


The groups R2 to R5 are selected from OH, OR and N(Y)Z such that:

At least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are OH,

15

At least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are OR, where R is defined above, with the proviso that when two of the groups R2 to R5 are OR, the R groups may not both be methyl or unsubstituted benzyl,

At least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are N(Y)Z, where Z is selected from hydrogen or R and Y is selected from the following, where G denotes the point of connection to the nitrogen atom in N(Y)Z, the N(Y)Z moieties may not be the same;
20



and the groups Q and W are independently selected from hydrogen or R as is defined above, and Q and W may combine to form a cycle,

- 5 The groups Z and Y may combine to form a cycle, and
The groups R1 to R5 may not combine together to form a cycle.

In a more particular form the invention resides in a compound as described above with the proviso that where two groups in the compound of formula I are N(Y)Z, these groups are different, with the further proviso that

10 when either R2 or R5 is N(Y)Z, N(Y)Z may not be azido, acetyl, benzyloxycarbonyl or t-butoxycarbonyl, with the further proviso that when R2 is N(Y)Z, N(Y)Z may not be phthalimido, 4-[N-[1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-3-methylbutyl]-amino]benzyl ester (ODmab), N-1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl (Dde), 2,2,2-Trichloroethoxycarbonyl (Troc), 9-


15 Fluorenylmethoxycarbonyl (Fmoc), or a 5-Acyl-1,3-dimethylbarbiturate type protecting group (DTPM) and with the further proviso that when the scaffold is of the 2-deoxy-2-aminoglucose configuration and R5 and R4 are both hydroxyl, R3 may not be a glycolate [$-CH_2-CO_2H$] or lactate ether [$-CH(CH_3)-CO_2H$] or an ester or amide derivative thereof.

Suitably, the compound is a derivative of a furanose form of a monosaccharide, and wherein n is 0.

Suitably, the compound is a derivative of a furanose form of a monosaccharide, and wherein n is 0.

- 5 Suitably, the compound has n = 1, at least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are N(Y)Z, where Z is selected from hydrogen or R and Y is selected from the following, where G denotes the point of connection to the nitrogen atom in N(Y)Z, the N(Y)Z moieties may not be the same;

10



- And the groups Q and W are independently selected from hydrogen or R as is defined above, with the proviso that Y and Z may not both
 15 be hydrogen and where two groups in the compound of formula I are N(Y)Z, these groups are different, the groups Z and Y may combine to form a cycle, the groups R1 to R5 may not combine together to form a cycle, with the proviso that where two groups in the compound of formula I are N(Y)Z, these groups are different, with the further proviso that when either R2 or R5 is N(Y)Z, N(Y)Z may
 20 not be azido, acetyl, benzyloxycarbonyl or t-butoxycarbonyl, with the further proviso that when R2 is N(Y)Z, N(Y)Z may not be phthalimido, 4-[N-[1-(4,4-

dimethyl-2,6-dioxocyclo-hexylidene)-3-methylbutyl]-amino}benzyl ester
(ODmab), *N*-1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl (Dde), 2,2,2-
Trichloroethoxycarbonyl (Troc), 9-Fluorenylmethoxycarbonyl (Fmoc), or a 5-
Acyl-1,3-dimethylbarbiturate type protecting group (DTPM) with the further
5 proviso that when the scaffold is of the 2-deoxy-2-aminoglucose configuration
and R5 and R4 are both hydroxyl, R3 may not be a glycolate [-CH₂-CO₂H] or
lactate ether [-CH(CH₃)-CO₂H] or an ester or amide derivative thereof.

Suitably the heteroarylalkyl is substituted by a moiety from the
group consisting of OH, NO, NO₂, NH₂, N₃, halogen, CF₃, CHF₂, CH₂F, nitrile,
10 alkoxy, aryloxy, amidine, guanidiniums, carboxylic acid, carboxylic acid ester,
carboxylic acid amide, aryl, cycloalkyl, heteroalkyl, heteroaryl, aminoalkyl,
aminodialkyl, aminotrialkyl, aminoacyl, carbonyl, substituted or unsubstituted
imine, sulfate, sulfonamide, phosphate, phosphoramidate, hydrazide,
hydroxamate, hydroxamic acid, heteroaryloxy, aminoalkyl, aminoaryl;
15 aminoheteroaryl, thioalkyl, thioaryl or thioheteroaryl, which may be further
substituted, with the proviso that the group R may not be or contain another
saccharide moiety, a peptide, protein or amino acid.

The compound may be immobilized to a support. The support
may be soluble or insoluble. Non-limiting examples of insoluble supports include
20 derivatised polystyrene, tentagel, wang resin, MBHA resin,
aminomethylpolystyrene, rink amide resin etc. Non-limiting examples of soluble
supports include DOX-mpeg, polyethylene glycol etc.

DETAILED DESCRIPTION

Embodiments of the invention will be described with reference to
25 the following examples. Where appropriate, the following abbreviations are
used.

| | |
|-------|--------------------------------|
| Ac | Acetyl |
| DTPM | 5-Acyl-1,3-dimethylbarbiturate |
| 30 Ph | Phenyl |
| TBDMS | <i>t</i> -Butyldimethylsilyl |
| TBDPS | <i>t</i> -Butyldiphenylsilyl |

| | | |
|----|--------|------------------------------------------------------------|
| | Bn | benzyl |
| | Bz | benzoyl |
| | Me | methyl |
| | DCE | 1,2-dichloroethane |
| 5 | DCM | dichloromethane, methylene chloride |
| | Tf | trifluoromethanesulfonyl |
| | Ts | 4-methylphenylsulfonyl, p-toluenesulfonyl |
| | DMF | N,N-dimethylformamide |
| | DMAP | N,N-dimethylaminopyridine |
| 10 | □□-DMT | □□-dimethoxytoluene, benzaldehyde dimethyl acetal |
| | DMSO | dimethylsulfoxide |
| | DTT | dithiothreitol |
| | DMTST | Dimethyl(methylthio)sulphoniumtrifluoro- methanesulphonate |
| | TBAF | tetra-n-butylammonium fluoride |


15

Part A: Preparation of building blocks:

In order to fully enable the invention, we detail below methods for the preparation of certain building blocks used in the preparation of the compounds of the invention. The building blocks described are suitable for both solution and solid phase synthesis of the compounds of the invention.


Example A: Synthesis of a 2,4 dinitrogen containing Galactopyranoside Building Block

12



Conditions: (i) 1,2-dimethoxytoluene (1,2-DMT), *p*-toluenesulphonic acid (TsOH), acetonitrile (MeCN), 76°C, 85%; (ii) Benzoylchloride (BzCl), triethylamine; DCM, 99%; (iii) methanol (MeOH)/MeCN/water, TsOH, 75°C, 98%; (iv) *t*-butyldiphenylsilylchloride (TBDPS-Cl), imidazole, pyridine, 120°C, 99%; (v) Tf₂O, pyridine, DCM, 0°C, 100%; (b) NaN₃, DMF, 16hr, RT, 99%.

Example B: Synthesis of a 3-nitrogen containing Gulopyranoside Building Block




5

Conditions: (i) (a) trifluoromethanesulfonic anhydride (Tf₂O), pyridine, -20°C, dichloromethane (DCM), 1 hour, 100%, (b) sodium azide (NaN₃), N,N-dimethylformamide (DMF), 50°C, 5 hours, quantitative; (ii) TsOH, MeCN/MeOH/water (12:3:1), 90°C, 6 hours, 88% (iii) TBDPSCI, DMAP, pyridine,


10 120°C, 12 hours, 93%

Example C: Synthesis of a 2,6-dinitrogen substituted Glucopyranoside Building Block



15


Conditions: (i) (a) Tosylchloride, pyridine, RT, 24 hours, 33% (b) NaN₃, DMF, RT, 168 hours.

Example D: Synthesis of a 2-nitrogen containing Tallopyranoside Building Block

5

Conditions: (i) TBDPSCl, imidazole, 1,2-DCE, reflux; (ii) NaOMe/MeOH; (iii) (a) TiF_2O , pyridine, -20°C , DCM, 1 hour, (b) NaN_3 , DMF, 50°C , 5 hours; (iv) TsOH, MeCN/MeOH/water; (v) benzoylchloride, DMAP, 1,2-DCE, -20°C .


Example E: Synthesis of two 3-nitrogen containing Altopyranoside Building Block




5

Conditions: (i) cyclohexanone dimethylacetal, TsOH, MeCN; (ii) *p*-methoxybenzaldehyde dimethylacetal, TsOH, MeCN; (iii) DIBAL, -78°C, diethyl ether; (iv) (a) Tf₂O, pyridine, -20°C, DCM, 1 hour, (b) NaN₃, DMF, 50°C, 5 hours; (v) TsOH, MeCN/MeOH/water; (vi) TBDPSCI, DMAP, 1,2-DCE; (vii) (a) CAN, (b) BzCl, DMAP, 1,2-DCE, (c) TsOH, MeCN/MeOH/water; (viii) TBDPSCI, DMAP, 1,2-DCE.

Example F: Synthesis of a 2-nitrogen containing Glucopyranoside Building Block




- 5 Conditions: (i) Ph-DMT, TsOH, MeCN; (ii) 1,2-DCE, BzCl, DMAP; (iii) TsOH, MeOH/MeCN; (iv) TBDPS-Cl, DMAP, 1,2-DCE.




- Conditions: (i) TBDPS-Cl, DMAP, pyridine, 120°C, 0.5 hours, 81%; (ii) a. (Bu)₂SnO, MeOH; b. Benzoylchloride, RT, 24 hour;

Example G: Synthesis of a 2-nitrogen containing Allopyranoside Building Block




- 5 *Conditions:* (i) DCM/pyridine, MsCl, DMAP, 0°C; (ii) sodium benzoate, dimethylsulphoxide (DMSO), 140°C; (iii) TsOH, MeOH/MeCN/water; (iv) TBDPS-Cl, imidazole, DCM, 1 hour, reflux.

Example H: Synthesis of a 3-nitrogen containing Allopyranoside Building Block


- 5 **Conditions:** (i) TiF_2O , pyridine, DCM; (b) NaN_3 , DMF; (ii) acetone, H^+ ; (iii) Ac_2O , pyridine; (iv) hexamethyldisilazane, I_2 , $\text{CH}_3\text{-S-S-CH}_3$; (v) NaOMe/MeOH ; (vi) TsOH , 1,4-dimethoxytoluene, MeCN; (vii) benzoylchloride, 1,2-DCE, pyridine, DMAP; (viii) TsOH , MeOH, H_2O , MeCN; (ix) TBDPS-Cl, imidazole, 1,2-DCE.

Example I: Syntheses of two 2-nitrogen containing Tallopyranoside Building Blocks with hydroxyls in the 3 or 4 positions.


5



Conditions: (i) (a) $\text{ Tf}_2\text{O/Py}$, (b) NaN_3 , DMF; (ii) TsOH , MeOH/MeCN/water ; (iii)
 10 BzCl , DMAP, 1,2-DCE; (iv) (a) phenoxyacetyl-Cl (PACl)/pyridine; (b)
 $\text{Bz}_2\text{O/pyridine}$; (v) MeNH_2/THF .

Example J: Synthesis of nitrogen containing furanoside Building Blocks

Conditions: (i) (a). 2,2-dimethoxypropane, TsOH, DMF; (b). TBDPSi-Cl, Imidazole, DMF; (ii) (a) $\text{TiF}_2\text{O}/\text{Py}$, (b) NaN_3 , DMF; (iii) (a) TsOH, MeOH/MeCN/water; (b) Benzoyl chloride, pyridine, DCM; (iv) 4-methoxybenzyl chloride, NaH, DMF; (v) (a) TBAF, THF; (b) $\text{TiF}_2\text{O}/\text{Py}$, (c) NaN_3 , DMF; (d) TsOH, MeOH/MeCN/water; (e) Benzoyl chloride, pyridine, DCM; (vi) (a) TsOH, MeOH/MeCN/water; (b) Benzoyl chloride, pyridine, DCM; (c) R-OH or R-SH, boron trifluoride diethyl etherate, DCM, molecular sieves; (d) $\text{TiF}_2\text{O}/\text{Py}$, (e) NaN_3 , DMF;

Example K: Synthesis of a 3-nitrogen containing Gulopyranoside Building Block


- Conditions:* (i) (a) trifluoromethanesulfonic anhydride (Tf_2O), pyridine, -20°C ,
 5 dichloromethane (DCM), 1 hour, 100%, (b) sodium azide (NaN_3), *N,N*-
 dimethylformamide (DMF), 50°C , 5 hours, quantitative; (ii)
 NaOH/ H_2O /THF/MeOH, 99%; (iii) Levulinic acid, *N,N'*-dicyclohexyldiimide,
 DMAP, DCM, quantitative; (iv) TsOH, MeCN/ MeOH/water (15:15:1), 50°C , 16
 hours, 56%; (v) TBDPSCI, DMAP, pyridine, 120°C , 2 hours, 85%; (vi)
 10 Benzoylchloride, pyridine, RT, 2 hour, 95%; (vii) hydrazine acetate, DCM.

Part B: Immobilization to solid support and glycosylation:

- The compounds of the present invention may be conveniently prepared in
 15 solution phase or on a solid support. Because a free hydroxyl group is always
 present in the compounds of the invention, it is convenient to immobilize the

building blocks to the solid support through a hydroxy function which will become the free hydroxyl group in the final compounds. Many of the building blocks described above have a free hydroxyl in the 4 position which is suitable for immobilization. Where a free hydroxyl is desired in a different position, a protection/deprotection sequence is first performed.

Example L: Alternative immobilization positions



10 **Conditions:** (i) 4-methoxybenzyl chloride, NaH, DMF, workup with citric acid (ii) NaOMe/MeOH/THF; (iii) TBAF/THF; HOAc to neutral pH

Example M: Glycosylation of anomeric position

15 In most cases the thiomethyl glycoside building block containing one free hydroxyl group can be used in glycosylation reactions without resorting to protection of the free hydroxyl. An excess of the alcohol acceptor is typically employed. Where a thiol is to be glycosylated, the acceptor alcohol is in short supply or results are not satisfactory, the thiomethyl glycoside donor may first

20 be converted to the bromo sugar or imidate, and these donors used for glycosylation. Alternatively, glycosylation can be effected with the fully protected precursor e.g. K-2, if significant side reaction is observed with the free hydroxy donors e.g. K-3, K-4, G-4.

In a typical procedure, 1 mmol of donor (eg G-4, K-2, K-3, K-4, A-6, B-4, C-1 etc) is dissolved in anhydrous dichloromethane 8 mL and an equal weight of dry 4A molecular sieves is added. The mixture is stirred for 30 minutes at room temperature then 4 mmol of the acceptor alcohol is added followed by addition
5 of DMTST solution (6 equivalents in 12ml of DCM). The reaction is monitored by t.l.c. When the reaction is complete, triethylamine (1.2 mmol) is added. The mixture is diluted with 100 mL dichloromethane and extracted with sodium bicarbonate (10% aqueous), citric acid (10% aqueous) and sodium chloride (sat. solution), dried over magnesium sulfate and solvents removed in vacuo.
10 The crude material is chromatographed on silica gel prior to immobilisation or in the case of K-2 removal of one of the alcohol protecting groups.

In an alternative procedure, 1 mmol of donor in dichloromethane 8 mL is first treated with bromine to yield the crude sugar halide. This solution is washed
15 briefly with 5% sodium thiosulfate, dried over magnesium sulfate and the solvents removed in vacuo. The crude sugar halide is used directly as above with silver triflate as the activating agent in place of DMTST. Both alcohols and thiols are amenable to glycosylation by this method.

20 Example N: Immobilization onto solid phase

Wang resin (13.3 g; 0.85 mmol/g, p-Benzyloxybenzyl Alcohol polystyrene-divinylbenzene resin) was dried in the vacuum oven overnight in 500 ml round bottom flask. The flask was place under nitrogen atmosphere then dry DCM
25 (133 ml) and trichloroacetonitrile (20 ml) was added. The mixture was cooled with ice bath while gently stirred. After 15 minutes of cooling DBU (1.3 ml) was added drop wise in 15 minutes, the resulting mixture was stirred for one hour with ice bath cooling. The resin was collected by filtering, washed with DMF, THF and DCM (3x each). The resin was dried in the vacuum oven over P₂O₅ for
30 24 hours to afford 15 grams of TriChloroAcetimidate Wang (TCA-Wang) resin. The resin was packed under nitrogen and stored at 4°C.
Yield 100%; loading ca. 0.754 mmol/g.

(Alternative resins may be used).

Glycosylated building blocks containing one free hydroxyl are immobilised onto TCA-Wang resin. In a typical procedure, TCA Wang resin (3.6 gram) was dried
5 in vacuum oven overnight then washed with anhydrous THF (3x36 ml) under nitrogen atmosphere. Building block (3 equiv.) was added followed by addition of anhydrous DCM (18 ml). The reaction mixture was shaken for 5 minutes (until all alcohol was dissolved), and $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.35 ml, 1 equivalent) was added. The reaction mixture was shaken vigorously for ten minutes and
10 drained; the resin was washed with DCM (3x30 ml), DMF (3x30 ml), THF (3x30 ml) and dried.

Part C: Library preparation:

The compounds of the invention are prepared by sequential deprotection and
15 ligation chemistries either on solid support or in solution phase. The following typical chemistries may be employed as required.

Removal of a tert-butyldiphenylsilyl:

The resin bound building block is suspended in dry THF/methanol (20/1 v/v)
20 mixture containing 10 equivalents of tetra-n-butylammonium fluoride. The mixture is stirred at 65°C for 24 hours, drained; the resin is filtered, washed with dimethylformamide followed by THF and finally dichloromethane. In an alternative procedure, TBAF may be conveniently replaced by HF.pyridine and the reaction effected in plastic ware. The TBAF may also be replaced by
25 HF."proton sponge" complex with good results.

Removal of a benzoate, p-chlorobenzoate or other ester protecting group:

The resin bound building block is suspended in dry THF and methanol (3/1 v/v)
mixture and sodium methoxide (0.5 equivalents) is added. The mixture is
30 shaken for 24 hours, drained and re-treated with fresh reagents for further 24 hours. The resin is filtered, washed with dimethylformamide followed by THF and finally dichloromethane.

Removal of a p-methoxybenzyl group:

The resin bound building block is suspended in DCM and a small amount of water is added (approx 1%) followed by 2,3-dichloro-5,6-dicyanobenzoquinone (10 equivalents). The mixture is shaken for 3 hours drained and re-treated with
5 fresh reagent for a further 3 hours. The resin is filtered, washed with THF followed by methanol and finally dichloromethane.

Etherification of hydroxyl position:

Resin bound building block which has previously had a hydroxyl group deprotected is washed three times and then suspended in anhydrous DMF and
10 3 equivalents of potassium t-butoxide added (alternative bases may be employed), shaken and drained after 5 minutes followed by the alkylating agent (3 equivalents) in DMF. The mixture is shaken for 10 minutes, drained and re-treated twice more with fresh reagents as above. The resin is filtered, washed
15 with dimethylformamide followed by THF and finally dichloromethane.

Reduction of an azide:

The resin bound building block is suspended in dry DMF; 5 equivalents of DTT (1,4-dithio-DL-threitol) and 3 equivalents of potassium tert-butoxide (alternative
20 bases may be employed) are added. The mixture is agitated under nitrogen atmosphere for 24 hours, drained and the resin is washed with dimethylformamide followed by THF and finally dichloromethane.

Removal of a DTPM group:

25 The resin bound building block is suspended in DMF and hydrazine hydrate (50/1 v/v) mixture, agitated 2 hours, drained and the resin is washed with dimethylformamide followed by THF and finally dichloromethane

Amide formation:

30 A solution of a suitable carboxylic acid (10 equivalents) in dry DMF is treated with HBTU (10 equivalents) and di-isopropylethylamine (10 equivalents) and shaken for 5 minutes. This solution is then added to a suspension of Resin

bound building block, which has previously had an amine group deprotected in DMF and the mixture shaken for 30 minutes. After this time the resin is drained and treated once more with fresh reagent for 30 minutes. The resin is filtered, washed with DMF followed by methanol and finally dichloromethane. If desired, quantitative ninhydrin assay may be performed to determine that the reaction is complete. Alternative coupling systems including HOAT, EDC/NHS or anhydrides may be employed to similar effect.

Urea and thiourea formation:

Isocyanates and thioisocyanates may be purchased or prepared by reaction of the corresponding amine with triphosgene, diphosgene, phosgene or thiophosgene as appropriate according to standard procedures as outlined in "Organic Functional Group Preparation" Vol I, 2nd Ed., Sandler and Karo, Academic Press, ISBN:0-12-618601-4 pp 359 to 375.

Resin bound building block which has previously had an amine group deprotected is suspended in anhydrous THF and 2 equivalents of the isocyanate or thioisocyanate added, followed immediately by triethylamine (1 equivalent). The mixture is shaken for 2 hours and may be exothermic depending on the scale and reactivity of the isocyanate or thioisocyanate used, drained and re-treated with fresh reagents for a further 2 hours. The resin is filtered, washed with THF followed by methanol and finally dichloromethane.

Carbamate formation:

Chloroformates and imidoylformates may be purchased or prepared by reaction of the corresponding alcohol with phosgene or carbonylbisimidazole as appropriate according to standard procedures as outlined in "Organic Functional Group Preparation" Vol I, 2nd Ed., Sandler and Karo, Academic Press, ISBN:0-12-618601-4 pp 359 to 375.

Resin bound building block which has previously had an amine group deprotected is suspended in anhydrous THF and 2 equivalents of the

chloroformate or imidoformate added, followed immediately by triethylamine (1 equivalent). The mixture is shaken for 2 hours and may be exothermic depending on the scale and reactivity of the isocyanate or thioisocyanate used, drained and re-treated with fresh reagents for a further 2 hours. The resin is
5 filtered, washed with THF followed by methanol and finally dichloromethane.

Sulfonamide formation:

Resin bound building block which has previously had an amine group deprotected is suspended in anhydrous THF or DMF and 2 equivalents of the
10 sulfonyl chloride added, followed immediately by triethylamine (2 equivalent). The mixture is shaken for 2 hours, drained and re-treated with fresh reagents for a further 2 hours. The resin is filtered, washed with THF or DMF followed by methanol and finally dichloromethane.

15 Removal of Fmoc:

The resin bound building block is suspended in piperidine /DMF (1/4, v/v) mixture and stirred 1 hours, drained and repeated once more; the resin is filtered, washed with dimethylformamide followed by THF and finally
dichloromethane.

20

Guanidine formation:












The resin bound building block is suspended in dry DMF containing 3 equivalents of 3,5-dimethylpyrazolyl formamidine nitrate and 15 equivalents of DIPEA. The mixture is stirred at 65°C for 24 hours, drained; the resin is
25 filtered, washed with dimethylformamide followed by THF and finally dichloromethane.

Cleavage of resin bound product:





















The resin bound compound is suspended in dry DCM containing 20% TFA and
30 20% Et₃SiH. The mixture is stirred at RT for 3 hours and the aliquot was collected; the resin was washed with dry DCM and all the DCM solutions were

combined, evaporated to dryness under reduced vacuo to furnish the desired product.










Libraries of compounds of the invention have been prepared based on the following scaffolds:

Scaffold **W1** derived from A-6Scaffold **W2** derived from B-4 or K-8Scaffold **W3** derived from C-1Scaffold **W4** derived from D-5 or I-6Scaffold **W5** derived from E-9 & E-7Scaffold **W6** derived from F-5 or F-7Scaffold **W7** derived from G-4Scaffold **W8** derived from H-9Scaffold **W9** derived from J-4Scaffold **W10** derived from J-6Scaffold **W11** derived from J-7

The following groups are exemplary of moieties in position R1, where the wavy line indicates the point of attachment to the carbohydrate ring:



31




The following groups are exemplary of ether linked moieties, where the wavy line indicates the point of attachment to an oxygen on the carbohydrate ring:

Methyl


Y1

Ethyl


Y2




Y3




Y4




Y5




Y6




Y7




Y8




Y9




Y10




Y11




Y12




Y13




Y14




Y15




Y16




Y17



Y18



Y19



Y20


33

Octyl.


Y21

Y22


Y23




Y24




Y25




Y26




Y27



Y28




Y29




Y30


The following groups are exemplary of amine linked moieties, where the wavy line indicates the point of attachment to a nitrogen on the carbohydrate ring:




Z1




Z2




Z3




Z4




Z5




Z6




Z7




Z8




Z9




Z10




Z11




Z12




Z13




Z14




Z15




Z16




Z17




Z18




Z19




Z20




Z21




Z22




Z23




Z24




Z25




Z26




Z27




Z28




Z29




Z30



Z31




Z32




Z33


36




Z34




Z35




Z36




Z37




Z38




Z39



Z40




Z41




Z42


38




Z56




Z57



Z58



Z59



Z60

Exemplary library compounds:

| Compound Number | Scaffold | R1 | R2 | R3 | R4 | R5 |
|-----------------|----------|----|-----|----|----|-----|
| 1 | W6 | X1 | Z43 | Y3 | OH | Y21 |
| 2 | W6 | X1 | Z44 | Y3 | OH | Y22 |
| 3 | W6 | X1 | Z45 | Y3 | OH | Y23 |
| 4 | W6 | X1 | Z46 | Y3 | OH | Y24 |
| 5 | W6 | X1 | Z47 | Y3 | OH | Y25 |
| 6 | W6 | X1 | Z48 | Y3 | OH | Y26 |
| 7 | W6 | X1 | Z49 | Y3 | OH | Y27 |
| 8 | W6 | X1 | Z50 | Y3 | OH | Y28 |
| 9 | W6 | X1 | Z51 | Y3 | OH | Y29 |
| 10 | W6 | X1 | Z52 | Y3 | OH | Y30 |
| 11 | W6 | X1 | Z53 | Y3 | OH | Y21 |
| 12 | W6 | X1 | Z54 | Y3 | OH | Y22 |
| 13 | W6 | X1 | Z55 | Y3 | OH | Y23 |
| 14 | W6 | X1 | Z56 | Y3 | OH | Y24 |
| 15 | W6 | X1 | Z57 | Y3 | OH | Y25 |
| 16 | W6 | X1 | Z58 | Y3 | OH | Y26 |
| 17 | W6 | X1 | Z59 | Y3 | OH | Y27 |
| 18 | W6 | X1 | Z60 | Y3 | OH | Y28 |
| 19 | W6 | X3 | Z12 | Y9 | OH | Y29 |
| 20 | W6 | X3 | Z29 | Y9 | OH | Y30 |
| 21 | W6 | X3 | Z12 | Y9 | OH | Y12 |
| 22 | W6 | X3 | Z29 | Y9 | OH | Y12 |
| 23 | W6 | X3 | Z13 | Y9 | OH | Y8 |
| 24 | W6 | X3 | Z26 | Y9 | OH | Y8 |
| 25 | W6 | X3 | Z13 | Y3 | OH | Y10 |
| 26 | W6 | X3 | Z26 | Y3 | OH | Y10 |
| 27 | W6 | X4 | Z3 | Y3 | OH | Y8 |
| 28 | W6 | X4 | Z17 | Y3 | OH | Y8 |

| | | | | | | |
|----|----|----|-----|-----|----|-----|
| 29 | W6 | X4 | Z3 | Y3 | OH | Y10 |
| 30 | W6 | X4 | Z17 | Y3 | OH | Y10 |
| 31 | W6 | X4 | Z12 | Y3 | OH | Y9 |
| 32 | W6 | X4 | Z29 | Y3 | OH | Y9 |
| 33 | W6 | X4 | Z3 | Y12 | OH | Y8 |
| 34 | W6 | X4 | Z17 | Y12 | OH | Y8 |
| 35 | W6 | X4 | Z3 | Y12 | OH | Y10 |
| 36 | W6 | X4 | Z17 | Y12 | OH | Y10 |
| 37 | W6 | X4 | Z12 | Y12 | OH | Y9 |
| 38 | W6 | X4 | Z29 | Y12 | OH | Y9 |
| 39 | W6 | X4 | Z3 | Y8 | OH | Y3 |
| 40 | W6 | X4 | Z17 | Y8 | OH | Y3 |
| 41 | W6 | X4 | Z3 | Y8 | OH | Y12 |
| 42 | W6 | X4 | Z17 | Y8 | OH | Y12 |
| 43 | W6 | X4 | Z13 | Y8 | OH | Y9 |
| 44 | W6 | X4 | Z26 | Y8 | OH | Y9 |
| 45 | W6 | X4 | Z3 | Y10 | OH | Y3 |
| 46 | W6 | X4 | Z17 | Y10 | OH | Y3 |
| 47 | W6 | X4 | Z3 | Y10 | OH | Y12 |
| 48 | W6 | X4 | Z17 | Y10 | OH | Y12 |
| 49 | W6 | X4 | Z13 | Y10 | OH | Y9 |
| 50 | W6 | X4 | Z26 | Y10 | OH | Y9 |
| 51 | W6 | X4 | Z12 | Y9 | OH | Y3 |
| 52 | W6 | X4 | Z29 | Y9 | OH | Y3 |
| 53 | W6 | X4 | Z12 | Y9 | OH | Y12 |
| 54 | W6 | X4 | Z29 | Y9 | OH | Y12 |
| 55 | W6 | X4 | Z13 | Y9 | OH | Y9 |
| 56 | W6 | X4 | Z26 | Y9 | OH | Y9 |
| 57 | W6 | X4 | Z13 | Y9 | OH | Y10 |
| 58 | W6 | X4 | Z26 | Y9 | OH | Y10 |
| 59 | W6 | X4 | Z3 | Y2 | OH | Y8 |


| | | | | | | |
|----|----|-----|-----|-----|----|-----|
| 60 | W6 | X4 | Z17 | Y2 | OH | Y8 |
| 61 | W6 | X4 | Z3 | Y2 | OH | Y10 |
| 62 | W6 | X4 | Z17 | Y2 | OH | Y10 |
| 63 | W6 | X4 | Z12 | Y2 | OH | Y9 |
| 64 | W6 | X4 | Z29 | Y2 | OH | Y9 |
| 65 | W6 | X4 | Z3 | Y8 | OH | Y1 |
| 66 | W6 | X10 | Z17 | Y8 | OH | Y1 |
| 67 | W6 | X10 | Z3 | Y8 | OH | Y2 |
| 68 | W6 | X10 | Z17 | Y8 | OH | Y2 |
| 69 | W6 | X10 | Z1 | Y8 | OH | Y9 |
| 70 | W6 | X10 | Z4 | Y8 | OH | Y9 |
| 71 | W6 | X10 | Z3 | Y10 | OH | Y1 |
| 72 | W6 | X10 | Z17 | Y10 | OH | Y1 |
| 73 | W6 | X10 | Z3 | Y10 | OH | Y2 |
| 74 | W6 | X10 | Z17 | Y10 | OH | Y2 |
| 75 | W6 | X10 | Z1 | Y10 | OH | Y9 |
| 76 | W6 | X10 | Z4 | Y10 | OH | Y9 |
| 77 | W6 | X10 | Z12 | Y9 | OH | Y1 |
| 78 | W6 | X10 | Z29 | Y9 | OH | Y1 |
| 79 | W6 | X10 | Z12 | Y9 | OH | Y2 |
| 80 | W6 | X10 | Z29 | Y9 | OH | Y2 |
| 81 | W6 | X10 | Z1 | Y9 | OH | Y9 |
| 82 | W6 | X10 | Z4 | Y9 | OH | Y9 |
| 83 | W6 | X15 | Z11 | Y1 | OH | Y17 |
| 84 | W6 | X15 | Z4 | Y9 | OH | Y10 |
| 85 | W8 | X6 | Y8 | Z33 | OH | Y9 |
| 86 | W8 | X6 | Y10 | Z24 | OH | Y19 |
| 87 | W8 | X6 | Y7 | Z18 | OH | Y12 |
| 88 | W8 | X9 | Y9 | Z25 | OH | Y3 |
| 89 | W8 | X9 | Y19 | Z1 | OH | Y4 |
| 90 | W8 | X9 | Y12 | Z20 | OH | Y13 |


| | | | | | | |
|-----|----|-----|-----|-----|-----|-----|
| 91 | W8 | X12 | Y3 | Z25 | OH | Y17 |
| 92 | W8 | X12 | Y4 | Z20 | OH | Y11 |
| 93 | W8 | X12 | Y13 | Z20 | OH | Y18 |
| 94 | W8 | X10 | Y17 | Z36 | OH | Y8 |
| 95 | W8 | X10 | Y11 | Z42 | OH | Y10 |
| 96 | W8 | X10 | Y18 | Z18 | OH | Y13 |
| 97 | W1 | X6 | Z33 | Y4 | Z37 | OH |
| 98 | W1 | X6 | Z37 | OH | Z33 | Y3 |
| 99 | W1 | X6 | Z42 | OH | Z18 | Y3 |
| 100 | W1 | X9 | Z33 | Y4 | Z37 | OH |
| 101 | W1 | X9 | Z37 | OH | Z33 | Y3 |
| 102 | W1 | X9 | Z42 | OH | Z18 | Y3 |
| 103 | W1 | X12 | Z33 | Y4 | Z37 | OH |
| 104 | W1 | X12 | Z37 | OH | Z33 | Y3 |
| 105 | W1 | X12 | Z42 | OH | Z18 | Y3 |
| 106 | W6 | X12 | Z11 | Y5 | OH | Y1 |
| 107 | W6 | X12 | Z16 | Y5 | OH | Y1 |
| 108 | W6 | X12 | Z5 | Y5 | OH | Y1 |
| 109 | W6 | X12 | Z11 | Y17 | OH | Y1 |
| 110 | W6 | X12 | Z16 | Y17 | OH | Y1 |
| 111 | W6 | X12 | Z5 | Y17 | OH | Y1 |
| 112 | W6 | X12 | Z11 | Y3 | OH | Y1 |
| 113 | W6 | X12 | Z16 | Y3 | OH | Y1 |
| 114 | W6 | X12 | Z5 | Y3 | OH | Y1 |
| 115 | W6 | X12 | Z11 | Y4 | OH | Y1 |
| 116 | W6 | X12 | Z16 | Y4 | OH | Y1 |
| 117 | W6 | X12 | Z5 | Y4 | OH | Y1 |
| 118 | W6 | X9 | Z11 | Y5 | OH | Y1 |
| 119 | W6 | X9 | Z16 | Y5 | OH | Y1 |
| 120 | W6 | X9 | Z5 | Y5 | OH | Y1 |
| 121 | W6 | X9 | Z11 | Y17 | OH | Y1 |

| | | | | | | |
|-----|----|-----|-----|-----|----|-----|
| 122 | W6 | X9 | Z16 | Y17 | OH | Y1 |
| 123 | W6 | X9 | Z5 | Y17 | OH | Y1 |
| 124 | W6 | X9 | Z11 | Y3 | OH | Y1 |
| 125 | W6 | X9 | Z16 | Y3 | OH | Y1 |
| 126 | W6 | X9 | Z5 | Y3 | OH | Y1 |
| 127 | W6 | X9 | Z11 | Y4 | OH | Y1 |
| 128 | W6 | X9 | Z16 | Y4 | OH | Y1 |
| 129 | W6 | X9 | Z5 | Y4 | OH | Y1 |
| 130 | W6 | X12 | Z11 | Y1 | OH | Y5 |
| 131 | W6 | X12 | Z16 | Y1 | OH | Y5 |
| 132 | W6 | X12 | Z5 | Y1 | OH | Y5 |
| 133 | W6 | X19 | Z28 | Y1 | OH | Y3 |
| 134 | W6 | X19 | Z13 | Y1 | OH | Y17 |
| 135 | W6 | X19 | Z13 | Y17 | OH | Y1 |
| 136 | W6 | X3 | Z29 | Y12 | OH | Y9 |
| 137 | W6 | X3 | Z17 | Y8 | OH | Y3 |
| 138 | W6 | X3 | Z17 | Y8 | OH | Y12 |
| 139 | W7 | X12 | Z11 | Y11 | OH | Y1 |
| 140 | W7 | X12 | Z16 | Y15 | OH | Y1 |
| 141 | W7 | X12 | Z3 | Y16 | OH | Y1 |
| 142 | W7 | X8 | Z11 | Y11 | OH | Y1 |
| 143 | W7 | X8 | Z16 | Y15 | OH | Y1 |
| 145 | W7 | X8 | Z3 | Y16 | OH | Y1 |
| 146 | W7 | X15 | Z11 | Y11 | OH | Y1 |
| 147 | W7 | X15 | Z16 | Y15 | OH | Y1 |
| 148 | W7 | X15 | Z3 | Y16 | OH | Y1 |
| 149 | W7 | X17 | Z11 | Y4 | OH | Y1 |
| 150 | W7 | X15 | Z7 | OH | Y4 | Y17 |
| 151 | W7 | X15 | Z31 | OH | Y4 | Y17 |
| 152 | W7 | X15 | Z9 | OH | Y4 | Y17 |
| 153 | W7 | X15 | Z32 | OH | Y4 | Y17 |

| | | | | | | |
|-----|-----|-----|-----|-----|--------|-----|
| 154 | W6 | X15 | Z42 | Y6 | Y1 | OH |
| 155 | W6 | X15 | Z37 | Y20 | Y1 | OH |
| 156 | W6 | X15 | Z39 | Y2 | Y1 | OH |
| 157 | W6 | X14 | Z42 | Y6 | Y8 | OH |
| 158 | W6 | X14 | Z37 | Y20 | Y8 | OH |
| 159 | W6 | X6 | Z17 | Y8 | Y3 | OH |
| 160 | W2 | X8 | OH | Z13 | Y4 | Y1 |
| 161 | W2 | X8 | OH | Z16 | Y4 | Y1 |
| 162 | W3 | X15 | Z36 | Y4 | OH | Z37 |
| 163 | W3 | X5 | Z11 | Y4 | OH | Z33 |
| 164 | W3 | X5 | Z8 | Y4 | OH | Z24 |
| 165 | W3 | X5 | Z36 | Y4 | OH | Z37 |
| 166 | W3 | X1 | Z11 | OH | OH | Z33 |
| 167 | W3 | X1 | Z8 | OH | OH | Z24 |
| 168 | W3 | X1 | Z36 | OH | OH | Z37 |
| 169 | W3 | X15 | Z11 | Y4 | OH | Z33 |
| 170 | W3 | X15 | Z8 | Y4 | OH | Z24 |
| 171 | W4 | X12 | Z10 | Y4 | Y8 | OH |
| 172 | W4 | X12 | Z41 | Y8 | Y3 | OH |
| 173 | W5 | X8 | Y17 | Z13 | Y4 | OH |
| 174 | W5 | X8 | Y17 | Z16 | Y4 | OH |
| 175 | W9 | X22 | Y4 | Z3 | Absent | OH |
| 176 | W9 | X23 | Y5 | Z11 | Absent | OH |
| 177 | W9 | X26 | Y8 | Z3 | Absent | OH |
| 178 | W9 | X21 | Y17 | Z11 | Absent | OH |
| 179 | W10 | X3 | Y6 | OH | Absent | Z25 |
| 180 | W10 | X5 | Y12 | OH | Absent | Z30 |
| 181 | W10 | X10 | Y19 | OH | Absent | Z40 |
| 182 | W11 | X6 | Z25 | OH | Absent | Y6 |
| 183 | W11 | X8 | Z30 | OH | Absent | Y12 |
| 184 | W11 | X10 | Z40 | OH | Absent | Y19 |

Exemplary synthesis of compound 85 (W6-X15-Z11-Y1-OH-Y17) on solid phase.






Conditions: (i) a. Br_2 , DCM; b. 4-Chlorobenzylalcohol, AgOTf, DCM; (ii) TCA-Wang resin, $\text{BF}_3 \cdot \text{Et}_2\text{O}$, DCM, THF; (iii) NaOMe, THF, MeOH; (iv) a. KOBU^t , DMF; b. iodomethane, DMF; (v) HF.'proton sponge', AcOH, DMF, 65°C ; (vi) a. KOBU^t , DMF; b. 2-bromomethyl-naphthalene, DMF; (vii) 1,4-Dithio-DL-threitol, KOBU^t , DMF; (viii) HBTU, Fmoc-Gly-OH, DIPEA, DMF; (ix) piperidine/ DMF (1/4); (x) 3,5-dimethylpyrazolyl formamidineium nitrate, DIPEA, DMF; (xi) TFA, Et_3SiH , DCM.

LCMS method:

| | Time | water% | acetonitrile% | Flow (ml/min) |
|---|-------|--------|---------------|---------------|
| | 0.00 | 95.0 | 5.0 | 2.000 |
| | 1.00 | 95.0 | 5.0 | 2.000 |
| 5 | 7.00 | 0.0 | 100.0 | 2.000 |
| | 12.00 | 0.0 | 100.0 | 2.000 |

M+H = 557.3; Rt = 3.98 min

Exemplary synthesis of compound 159 (W6-Z17-Y8-Y3-OH) in solution phase:


- 5 Conditions: (i) 4-Methoxybenzaldehyde dimethylacetal, TsOH, CH₃CN; (ii) NaH (95%), tert-butyl bromoacetate, DMF; (iii) NaBH₃CN, TFA, DMF; (iv) KOBu^t, BnBr, DMF; (v) a. Zn, NH₄Cl, MeOH, H₂O; b. HBTU, 3-Boc-NH-benzoic acid, DIPEA, DMF; (vi) CH₃CN, H₂O, TsOH.

- 10 It should be appreciated that various changes and modifications can be made to the embodiments without departing from the spirit and scope of the invention.

CLAIMS:

1. A compound of formula I being a derivative of a furanose or pyranose form of a monosaccharide,

5



formula I

Wherein, n is 0 or 1;

R1 is XR wherein,

10

X is selected from O; S; S=O and SO₂,

R is selected from the group consisting of C1 to C9 alkyl, C1 to C15 alkenyl, C1 to C15 alkynyl, C1 to C15 heteroalkyl, C6 to C15 aryl, C6 to C15 heteroaryl, C6 to C15 arylalkyl or C6 to C15 heteroarylalkyl which is optionally substituted, cyclic or acyclic, branched and/or linear,


15

the groups R2 to R5 are selected from OH, OR and N(Y)Z such that:

at least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are OH,

20 at least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are OR, where R is defined above, with the proviso that when two of the groups R2 to R5 are OR, the R groups may not both be methyl or unsubstituted benzyl,

25 at least one of the groups R2 to R5 and not more than two of the groups R2 to R5 are N(Y)Z, where Z is selected from hydrogen or R and Y is selected from the following, where G denotes the point of connection to the nitrogen atom in N(Y)Z, the N(Y)Z moieties may not be the same;



and the groups Q and W are independently selected from hydrogen or R as is
 5 defined above, and Q and W may combine to form a cycle,
 the groups Z and Y may combine to form a cycle,
 the groups R1 to R5 may not combine together to form a cycle,

with the proviso that where two groups in the compound of formula
 I are N(Y)Z, these groups are different,

10 with the further proviso that when either R2 or R5 is N(Y)Z, N(Y)Z
 may not be azido, acetyl, benzyloxycarbonyl or t-butoxycarbonyl,

with the further proviso that when R2 is N(Y)Z, N(Y)Z may not be
 phthalimido, 4-[N-[1-(4,4-dimethyl-2,6-dioxocyclo-hexylidene)-3-methylbutyl]-
 amino}benzyl ester (ODmab), N-1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl
 15 (Dde), 2,2,2-Trichloroethoxycarbonyl (Troc), 9-Fluorenylmethoxycarbonyl
 (Fmoc), or a 5-Acyl-1,3-dimethylbarbiturate type protecting group (DTPM),

with the further proviso that when the scaffold is of the 2-deoxy-2-aminoglucose
 configuration and R5 and R4 are both hydroxyl, R3 may not be a glycolate [-


CH₂-CO₂H] or lactate ether [-CH(CH₃)-CO₂H] or an ester or amide derivative thereof.

2. The compound of claim 1 which is a derivative of a pyranose form of a monosaccharide and wherein n is 1.

3. The compound of claim 1 which is a derivative of a furanose form of a monosaccharide, and wherein n is 0.

4. The compound of claim 2, wherein n is 1,

at least one of the groups R₂ to R₅ and not more than two of the groups R₂ to R₅ are N(Y)Z, where Z is selected from hydrogen or R and Y is selected from the following, where G denotes the point of connection to the nitrogen atom in N(Y)Z, the N(Y)Z moieties may not be the same;




and the groups Q and W are independently selected from hydrogen or R as is defined above, with the proviso that Y and Z may not both be hydrogen and


- where two groups in the compound of formula I are N(Y)Z, these groups are different,
- the groups Z and Y may combine to form a cycle,
- the groups R1 to R5 may not combine together to form a cycle,
- 5 with the proviso that where two groups⁹ in the compound of formula I are N(Y)Z, these groups are different,
- with the further proviso that when either R2 or R5 is N(Y)Z, N(Y)Z may not be azido, acetyl, benzyloxycarbonyl or t-butoxycarbonyl,
- with the further proviso that when R2 is N(Y)Z, N(Y)Z may not be phthalimido,
- 10 4-[N-[1-(4,4-dimethyl-2,6-dioxocyclo-hexylidene)-3-methylbutyl]-amino}benzyl ester (ODmab), *N*-1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl (Dde), 2,2,2-Trichloroethoxycarbonyl (Troc), 9-Fluorenylmethoxycarbonyl (Fmoc), or a 5-Acyl-1,3-dimethylbarbiturate type protecting group (DTPM),
- with the further proviso that when the scaffold is of the 2-deoxy-2-aminoglucose
- 15 configuration and R5 and R4 are both hydroxyl, R3 may not be a glycolate [$-\text{CH}_2\text{-CO}_2\text{H}$] or lactate ether [$-\text{CH}(\text{CH}_3)\text{-CO}_2\text{H}$] or an ester or amide derivative thereof.

5. The compound of any one of claim 1-4 wherein the heteroarylalkyl
- 20 is substituted by a moiety from the group consisting of OH, NO, NO₂, NH₂, N₃, halogen, CF₃, CHF₂, CH₂F, nitrile, alkoxy, aryloxy, amidine, guanidiniums, carboxylic acid, carboxylic acid ester, carboxylic acid amide, aryl, cycloalkyl, heteroalkyl, heteroaryl, aminoalkyl, aminodialkyl, aminotrialkyl, aminoacyl, carbonyl, substituted or unsubstituted imine, sulfate, sulfonamide, phosphate,
- 25 phosphoramidate, hydrazide, hydroxamate, hydroxamic acid, heteroaryloxy, aminoalkyl, aminoaryl, aminoheteroaryl, thioalkyl, thioaryl or thioheteroaryl, which may be further substituted, with the proviso that the group R may not be or contain another saccharide moiety, a peptide, protein or amino acid.


6. The compound of claim 1 which comprises, as a precursor




7. The compound of claim 1, which comprises as a precursor




- 5 8. The compound of claim 1, which comprises as a precursor



9. The compound of claim 1, which comprises as a precursor




10. The compound of claim 1, which comprises as a precursor




10


11. The compound of claim 1, which comprises as a precursor




12. The compound of claim 1, which comprises as a precursor




13. The compound of claim 1, which comprises as a precursor




- 5 14. The compound of claim 1, which comprises as a precursor



15. The compound of claim 1, which comprises as a precursor




16. The compound of claim 1, which comprises as a precursor




10


17. The compound of claim 1, which comprises as a precursor




18. The compound of claim 1, which comprises as a precursor




19. The compound of claim 1, which comprises as a precursor



20. The compound of claim 1, which comprises as a precursor



21. The compound of claim 1, which comprises as a precursor



22. The compound of claim 1 which is immobilised to a support.





















10










23. The compound of claim 22, wherein the compound is immobilised to the support through a hydroxyl group.

- 24 The compound of claim 23, wherein the support is selected from the group consisting of derivatised polystyrene, tentagel, wang resin, MBHA resin, aminomethylpolystyrene, rink amide resin, DOX-mpeg and polyethylene glycol.

15

25. The compound of claim 1, wherein R1 is selected from the group consisting of






26. The compound of claim 1, wherein one of the R moieties in OR is selected from the group consisting of

Methyl


Y1

Ethyl


Y2




Y3




Y4




Y5




Y6




Y7




Y8




Y9




Y10




Y11




Y12




Y13




Y14




Y15




Y16




Y17



Y18



Y19




Y20


59

Octyl


Y21




Y22




Y23




Y24




Y25




Y26




Y27



Y28





















Y29




Y30


27. The compound of claim 1, wherein Z is selected from the group consisting of




61




Z19




Z20




Z21




Z22




Z23




Z24




Z25




Z26




Z27




Z28




Z29




Z30



Z31




Z32




Z33


62




Z34




Z35




Z36




Z37




Z38




Z39




Z40




Z41




Z42




Z43




Z44




Z45




Z46




Z47




Z48




Z49




Z50




Z51




Z52




Z53




Z54


**Z55**




Z56




Z57



Z58




Z59



Z60

AMENDED CLAIMS

[received by the International Bureau on 07 November 2003 (07.11.03) ;
original claim 1 replaced by new claim 1 ;
remaining claims unchanged (1 page)]



and the groups Q and W are independently selected from hydrogen or R as is defined above, and Q and W may combine to form a cycle,

the groups Z and Y may combine to form a cycle,

the groups R1 to R5 may not combine together to form a cycle,

with the proviso that where two groups in the compound of formula I are N(Y)Z, these groups are different,


with the further proviso that when either R2 or R5 is N(Y)Z, N(Y)Z may not be trifluoroacetamido, acetamido, benzyloxycarbonylamino or t-butoxycarbonylamino,

with the further proviso that when R2 is N(Y)Z, N(Y)Z may not be phthalimido, 4-[N-[1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-3-methylbutyl]-amino]benzyl ester (ODmab), N-1-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl (Dde), 2,2,2-Trichloroethoxycarbonyl (Troc), 9-Fluorenylmethoxycarbonyl (Fmoc), or a 5-Acyl-1,3-dimethylbarbiturate type protecting group (DTPM),

with the further proviso that when the scaffold is of the 2-deoxy-2-aminoglucose configuration and R5 and R4 are both hydroxyl, R3 may not be a glycolate [-

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU03/01008

| | | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| A. CLASSIFICATION OF SUBJECT MATTER | | |
| Int. Cl. ⁷ : C07H 15/18, 5/06, 23/00 | | |
| According to International Patent Classification (IPC) or to both national classification and IPC | | |
| B. FIELDS SEARCHED | | |
| Minimum documentation searched (classification system followed by classification symbols) | | |
| Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched | | |
| Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) STN substructure search | | |
| C. DOCUMENTS CONSIDERED TO BE RELEVANT | | |
| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| X | Derwent Abstract Accession No. 2002-371191/40 Class B03 US 2002028927-A1 (CHRIST W. J.) 7 March 2002. See whole document. | 1-5 |
| X | US 6,184,366 B1 (CHRIST) 6 February 2001 See Claims and Examples; Columns 17-18 | 1-5, 6-21 |
| X | WO 02/32915 A1 (ALCHEMIA PTY LTD) 25 April 2002 See whole document. | 1-5 |
| <input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex | | |
| <p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p> | | |
| Date of the actual completion of the international search 3 September 2003 | | Date of mailing of the international search report 9 SEP 2003 |
| Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaustalia.gov.au Facsimile No. (02) 6285 3929 | | Authorized officer  S.R. IDRUS Telephone No : (02) 6283 2659 |

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU03/01008

| C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT | | |
|-------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| X | EP 0099578 B1 (CIBA GEIGY AG) 1 February 1984 (& US 4,548,923 See column 6, formula III) | 1-5 |
| X | MALETIC M et. al., "Preparation of potential inhibitors of the mur-Pathway enzymes on solid support using an acetal linker", Bioorganic & Medicinal Chemistry Letters (2003), 13(6), 1125-1128. See page 1127 col. 2 Fig.2 | 1-5, 26 |
| X | FUKASE Y et. al., "New efficient route for synthesis of lipid A by using affinity separation", Synlett (2001), (11), 1693-1698. See Scheme 1 compound 3, Scheme 3 compounds 17, 18, and 20 | 1-5 |
| X | NAKAYAMA K et. al., "Novel peptidomimetics of the antifungal cyclic peptide Rhodopeptin: design of mimetics utilizing scaffolding methodology", Organic Letters. (2001), 3 (22), 3447-3450. See page 3448 Fig. 4 compounds 5 and 6 | 1-5 |
| X | YOSHIZAKI H et. al., "First total synthesis of the Re-type lipopolysaccharide", Angewandte Chemie, International Edition (2001), 40(8), 1475-1480. See page 1477 col. 2 compounds 16 and 17 | 1-5 |
| X | HANESSIAN S et. al., "Formation of 4-alkoxybenzylidene acetals on solid support and generation of functional diversity with carbohydrate scaffolds", Synlett (1999), (1), 102-104 See whole document. | 1-5, 22-24 |

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU03/01008

Box I Observations where certain claims were found unsearchable (Continuation of Item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos :
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos : 1-5 (in parts), 6-21
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Claims 1-5 provide for such a vast number of possible compounds by virtue of the permutations of the variables present that they have not been fully searched. Claims 6-21 are directed to compounds of the invention as defined in Claims 1-5 wherein one or more of the hydroxyl oxygens are protected

3. ☐ Claims Nos :
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

Box II Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.;

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/AU03/01008

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

| Patent Document Cited in Search Report | | | | Patent Family Member | | | |
|-------------------------------------------|----------|----|----------|----------------------|----------|----|----------|
| US | 6184366 | AU | 63802/96 | CA | 2223140 | CN | 1192216 |
| | | EP | 853627 | HU | 9802662 | IL | 122251 |
| | | JP | 11506793 | NO | 975644 | NZ | 312299 |
| | | RU | 2170738 | ZA | 9604666 | | |
| WO | 02/32915 | EP | 1326873 | | | | |
| EP | 0099578 | AT | 23536 | AU | 17218/83 | CA | 1243305 |
| | | CS | 8305484 | CS | 8308406 | DD | 210058 |
| | | DK | 3378/83 | ES | 524386 | FI | 832639 |
| | | GR | 79611 | IL | 69292 | JP | 59033296 |
| | | NO | 832691 | NZ | 205002 | PL | 243132 |
| | | PT | 77081 | SU | 1299516 | US | 4548923 |
| | | ZA | 8305357 | | | | |
| | | | | | | | |
| END OF ANNEX | | | | | | | |